

CIRCLE K NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid
Lil' Drug Store Products, Inc

Circle K™ Nighttime Cold & Flu

Drug Facts

<i>Active ingredients (in each 30 mL)</i>	<i>Purposes</i>
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - runny nose and sneezing
 - fever
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain

drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts.
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with the dosing cup provided. Do not use any other dosing device.

- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years of age: do not use
- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

Other information

- **each 30 mL contains:** potassium 5 mg
- **each 30 mL contains:** sodium 24 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

acesulfame potassium, alcohol, citric acid, FD&C blue #1, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?

Call toll free **1-877-507-6516 (M-F 8AM-4:30PM CST)**

Proudly distributed by Circle K Stores Inc.

PRINCIPAL DISPLAY PANEL - 237 mL Bottle Label

Compare to the Active Ingredients in Vicks® NyQuil®*

CIRCLE K™

Nighttime Cold & Flu

Acetaminophen 650 mg, Pain Reliever/Fever Reducer

Dextromethorphan HBr 30 mg, Cough Suppressant

Doxylamine Succinate 12.5 mg, Antihistamine

Relieves

- Aches - Sore Throat - Fever
- Sneezing - Runny Nose - Cough

10% Alcohol

Cherry

8 FL OZ (237mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Drug Facts

Active ingredients (in each 30 mL) Purposes

Acetaminophen 650 mg.....Pain reliever/fever reducer
 Dextromethorphan HBr 30 mg.....Cough suppressant
 Doxylamine succinate 12.5 mg.....Antihistamine

Uses

temporarily relieves these common cold/flu symptoms:
 • minor aches and pains • headache • sore throat
 • runny nose and sneezing • fever
 • cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
 If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

◀ **LIFT LABEL FOR CONTINUED DRUG FACTS**

Compare to the Active Ingredients in Vicks® NyQuil®*



Nighttime Cold & Flu

Acetaminophen 650 mg, Pain Reliever/Fever Reducer
 Dextromethorphan HBr 30 mg, Cough Suppressant
 Doxylamine Succinate 12.5 mg, Antihistamine

Relieves

- Aches - Sore Throat - Fever
- Sneezing - Runny Nose - Cough

10% Alcohol
Cherry

8 FL OZ (237mL)

Product manufactured for:
 Lill Drug Store Products, Inc.
 9300 Earhart Lane SW
 Cedar Rapids, IA 52404

Proudly distributed by Circle K Stores, Inc.

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil® are registered trademarks of The Procter & Gamble Company.

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

PLD-B427B LB006158
 57344L-US-11-17-A



Drug Facts (continued)

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease • glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. ▶

Drug Facts (continued)

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with the dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years of age: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- each 30 mL contains: potassium 5 mg
- each 30 mL contains: sodium 24 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients aceulfame potassium, alcohol, citric acid, FD&C blue #1, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?
 Call toll free 1-877-507-6516 (M-F 8AM-4:30PM CST)

PDP

TEMPORARILY RELIEVES ACHES, SORE THROAT, FEVER, SNEEZING, RUNNY NOSE, COUGH

Compare to the active ingredients in **Vicks® NyQuil® Cold & Flu***

CIRCLE K

nighttime cold & flu
cherry flavor, 10% alcohol

**Acetaminophen 650 mg,
Dextromethorphan HBr 30 mg, Doxylamine succinate 12.5 mg
Pain Reliever/Fever Reducer, Cough Suppressant,
Antihistamine**

8 fl oz (237 mL)

<p>TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.</p> <p>Drug Facts</p> <table border="1"> <thead> <tr> <th>Active ingredients (in each 30 mL)</th> <th>Purposes</th> </tr> </thead> <tbody> <tr> <td>Acetaminophen 650 mg</td> <td>Pain reliever/fever reducer</td> </tr> <tr> <td>Dextromethorphan HBr 30 mg</td> <td>Cough suppressant</td> </tr> <tr> <td>Doxylamine succinate 12.5 mg</td> <td>Antihistamine</td> </tr> </tbody> </table> <p>Uses temporarily relieves these common cold/flu symptoms:</p> <ul style="list-style-type: none"> • minor aches and pains • headache • sore throat • runny nose and sneezing • fever • cough due to minor throat and bronchial irritation <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> • more than 4000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> • skin reddening • blisters • rash <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional 	Active ingredients (in each 30 mL)	Purposes	Acetaminophen 650 mg	Pain reliever/fever reducer	Dextromethorphan HBr 30 mg	Cough suppressant	Doxylamine succinate 12.5 mg	Antihistamine	<p>TEMPORARILY RELIEVES ACHES, SORE THROAT, FEVER, SNEEZING, RUNNY NOSE, COUGH Compare to the active ingredients in Vicks® NyQuil® Cold & Flu*</p> <div style="border: 2px solid blue; border-radius: 50%; padding: 20px; width: 80%; margin: 0 auto;"> <p>CIRCLE K™</p> <p>nighttime cold & flu</p> <p>cherry flavor, 10% alcohol</p> <p>Acetaminophen 650 mg, Dextromethorphan HBr 30 mg, Doxylamine succinate 12.5 mg</p> <p>Pain Reliever/Fever Reducer, Cough Suppressant, Antihistamine</p> </div> <p>8 fl oz (237 mL)</p>	<p>Product manufactured for: Lif Drug Store Products, Inc. 9300 Earhart Lane SW Cedar Rapids, IA 52404</p> <p>Proudly distributed by Circle K Stores, Inc.</p> <p>*This product is not manufactured or distributed by The Procter & Gamble Company, distributor of Vicks® NyQuil® Cold & Flu.</p> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p>PARENTS: Learn about best medicines about www.StopMedicineAbuse.org</p> </div> <p>PLD-C427B LB007944 55344L-US-11-17-3</p>  <p>1 94283 65157 5</p>
Active ingredients (in each 30 mL)	Purposes									
Acetaminophen 650 mg	Pain reliever/fever reducer									
Dextromethorphan HBr 30 mg	Cough suppressant									
Doxylamine succinate 12.5 mg	Antihistamine									

<p>Drug Facts (continued)</p> <p>conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> • liver disease • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating due to an enlarged prostate gland <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> • taking sedatives or tranquilizers • taking the blood thinning drug warfarin <p>When using this product</p> <ul style="list-style-type: none"> • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> • pain or cough gets worse or lasts more than 7 days • new symptoms occur • fever gets worse or lasts more than 3 days • redness or swelling is present • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p>	<p>Drug Facts (continued)</p> <p>Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> • do not take more than directed (see Overdose warning) • do not take more than 4 doses in any 24-hour period • measure only with the dosing cup provided. Do not use any other dosing device. • mL = milliliter • keep dosing cup with product • adults and children 12 years and over: 30 mL every 6 hours • children under 12 years of age: do not use • when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing <p>Other information</p> <ul style="list-style-type: none"> • each 30 mL contains: potassium 5 mg • each 30 mL contains: sodium 24 mg • store between 20-25°C (68-77°F). Do not refrigerate. <p>Inactive ingredients acesulfame potassium, alcohol, citric acid, FD&C blue #1, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate</p> <p>Questions or comments? Call toll free 1-877-507-6516 (M-F 8AM-4:30PM CST)</p>
--	--

CIRCLE K NIGHTTIME COLD AND FLU			
acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-5534
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ALCOHOL (UNII: 3K9958V90M)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5534-4	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/03/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/03/2021	

CIRCLE K NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-5834
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ALCOHOL (UNII: 3K9958V90M)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5834-4	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/15/2019	09/15/2024

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/01/2014	09/15/2024

Labeler - Lil' Drug Store Products, Inc (093103646)

Revised: 12/2023

Lil' Drug Store Products, Inc