

COLD AND FLU RELIEF- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid
Walgreens

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - headache
 - fever
 - sore throat
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- adult take more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blistere
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if user is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause serious liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-

800-222-1222) right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms

Directions

- **do not take more than directed (see overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- mL = milliliter
- keep dosing cup with product

| | |
|---------------------------------------|---------------------|
| adults and children 12 years and over | 30 mL every 4 hours |
| children 6 to under 12 years | 15 mL every 4 hours |
| children 4 to under 6 years | ask a doctor |
| children under 4 years | do not use |

- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

Other information

- each 15 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, FD&C Yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu††

DAYTIME • NON-DROWSY

Cold & Flu

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER

DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT

PHENYLEPHRINE HCl 5 mg / NASAL CONGESTION

Multi-Symptom

- Relieves aches, fever & sore throat , cough & nasal congestant
- For ages 6 years & over
- Alcohol free
- Antihistamine free

FL OZ (mL)

††This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

Product Label

Drug Facts (continued)

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 days (adult) ■ fever gets worse, or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** **Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ **do not take more than directed** (see Overdose warning) ■ **do not take more than 4 doses in any 24-hour period** ■ **measure only with dosing cup provided.** Do not use any other dosing device. ■ mL = milliliter ■ **keep dosing cup with product**

| | |
|---------------------------------------|---------------------|
| adults and children 12 years and over | 30 mL every 4 hours |
| children 6 to under 12 years | 15 mL every 4 hours |
| children 4 to under 6 years | ask a doctor |
| children under 4 years | do not use |

■ **when using other Daytime or Nighttime products**, carefully read each label to ensure correct dosing

Other information ■ each 15 mL contains: sodium 12 mg ■ store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

†Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. ††This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

DISTRIBUTED BY: WALGREEN CO. 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED Walgreens.com ©2022 Walgreen Co.

ITEM 271558 W10369-0922-F
 3 11917 21864 9
W30R03022-F
REV-0922

PARENTS:
Learn about your medicine online
www.StopMedicineAcute.org
PLD-B3800 LB0008967

Walgreens

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu††

DAYTIME • NON-DROWSY Cold & Flu

ACETAMINOPHEN 325 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT
PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT

Multi-Symptom

- Relieves aches, fever & sore throat, cough & nasal congestion
- For ages 6 years & over
- Alcohol free
- Antihistamine free

NDC 0363-4661-12

Drug Facts

Active ingredients (in each 15 mL)

| | |
|---------------------------------|-----------------------------|
| Acetaminophen 325 mg..... | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg..... | Cough suppressant |
| Phenylephrine HCl 5 mg..... | Nasal decongestant |

Uses ■ temporarily relieves common cold and flu symptoms ■ minor aches and pains ■ headache ■ fever ■ sore throat ■ nasal congestion ■ cough due to minor throat and bronchial irritation

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if: ■ adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount ■ child takes more than 4 doses (15 mL each) in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ a sodium-restricted diet ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not exceed recommended dosage.

WALGREENS Daytime Non-Drowsy Cold & Flu

COLD AND FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid

| Product Information | | | |
|---------------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:0363-4661 |
| Route of Administration | ORAL | | |
| Active Ingredient/Active Moiety | | | |

| Ingredient Name | Basis of Strength | Strength |
|---|----------------------------------|--------------------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg in 15 mL |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg in 15 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg in 15 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SORBITOL (UNII: 506T60A25R) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | |
| SODIUM CITRATE (UNII: 1Q73Q2JULR) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:0363-4661-12 | 355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 07/30/2020 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 07/30/2020 | |

Labeler - Walgreens (008965063)