

**DAYTIME NIGHTTIME MULTI SYMPTOM COLD AND FLU RELIEF-  
acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine  
hcl  
CVS Pharmacy**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**CVS Pharmacy, Inc. Daytime Nighttime Cold & Flu Relief Drug Facts**

**Nighttime Cold & Flu  
Active ingredients (in each 30 mL)**

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

### **Other information**

- **each 30 mL contains:** sodium 38 mg
- store at 20-25°C (68-77°F)

### **Inactive ingredients**

alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

### **Questions or comments?**

**1-800-719-9260**

### **Daytime Cold & Flu**

#### **Active ingredients (in each 15 mL)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

### **Purpose**

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

## Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**When using this product do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- **each 15 mL contains:** sodium 7 mg
- store at 20-25°C (68-77°F)

### **Inactive ingredients**

butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

### **Questions or comments?**

**1-800-719-9260**

### **Principal Display Panel**

COMBO PACK

CVS HEALTH®

CVS HEALTH®

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu

Non-Drowsy

Daytime

ACETAMINOPHEN - Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr - Cough suppressant

PHENYLEPHRINE HCl - Nasal decongestant

Relieves: Aches, Fever, Sore throat, Cough, Nasal congestion

COLD & FLU

ALCOHOL & ANTIHISTAMINE FREE

TOTAL 24 FL OZ (1.5 PT) (710 mL)

CVS HEALTH®

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu

Multi-Symptom

Nighttime

ACETAMINOPHEN - Pain reliever; Fever reducer

DEXTROMETHORPHAN HBr - Cough suppressant

DOXYLAMINE SUCCINATE - Antihistamine

Relieves: Aches, Fever, Sore throat, Cough, Sneezing, Runny nose

COLD & FLU

Original Flavor

ALCOHOL 10%

TWO 12 FL OZ (355 mL) UNITS

 **CVS**Health.

**Non-Drowsy  
Daytime**

**Multi-Symptom  
Nighttime**

**COMBO PACK**

 **CVS**Health.

 **CVS**  
Health.

NDC 59779-567-02

**Multi-Symptom  
Nighttime**

**ACETAMINOPHEN-**  
Pain reliever; Fever reducer  
**DEXTROMETHORPHAN HBr-**  
Cough suppressant  
**DOXYLAMINE SUCCINATE-**  
Antihistamine

 **CVS**  
Health.

Compare to the  
active ingredients  
in Vicks® DayQuil®  
Cold & Flu\*

NDC 59779-606-40

**Non-Drowsy  
Daytime**

**ACETAMINOPHEN-**  
Pain reliever; Fever reducer  
**DEXTROMETHORPHAN HBr-**  
Cough suppressant  
**PHE NYLEPHRINE HCl-**  
Nasal decongestant

 **CVS**  
Health.

Compare to the  
active ingredients  
in Vicks® NyQuil®  
Cold & Flu\*

NDC 59779-335-40

**Multi-Symptom  
Nighttime**

**ACETAMINOPHEN-**  
Pain reliever; Fever reducer  
**DEXTROMETHORPHAN HBr-**  
Cough suppressant  
**DOXYLAMINE SUCCINATE-**  
Antihistamine

Relieves: Aches, Fever, Sore throat,  
Cough, Nasal congestion

**COLD & FLU**

ALCOHOL &  
ANTIHISTAMINE FREE

**TOTAL 24 FL OZ  
(1.5 pt) (710 mL)**

Relieves: Aches, Fever, Sore throat,  
Cough, Sneezing, Runny nose

**COLD & FLU**

Original Flavor  
ALCOHOL 10%

**TWO 12 FL OZ  
(355 mL) UNITS**

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**DO NOT USE IF PRINTED NECKBAND  
IS BROKEN OR MISSING**

#890847



PAPER PLASTIC PLASTIC  
BOX BOTTLE CUP

0 50428 26419 5

56702 17 C5

# Nighttime Cold & Flu

## Drug Facts

### Active ingredients (in each 30 mL)

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

### Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### Ask a doctor before use if you have

- liver disease
- glucocoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

### Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- take only as directed - see Overdose warning

only use the dose cup provided	do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

### Other information

- each 30 mL contains: sodium 38 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients** alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

**Questions or comments?** 1-800-719-9260

# Daytime Cold & Flu

## Drug Facts

### Active ingredients (in each 15 mL)

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

### Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

### Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

### When using this product do not use more than directed

### Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- take only as directed - see Overdose warning

only use the dose cup provided	do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

### Other information

- each 15 mL contains: sodium 7 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients** butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, methylol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

**Questions or comments?** 1-800-719-9260



## Non-Drowsy

# Daytime

**ACETAMINOPHEN-**  
Pain reliever; Fever reducer

**DEXTROMETHORPHAN HBr-**  
Cough suppressant

**PHENYLEPHRINE HCl-**  
Nasal decongestant

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2022 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS  
V-16430

100% money back  
guaranteed.  
CVS.com/returnpolicy

\*These products are not manufactured or distributed  
by Procter & Gamble, distributor of Vicks® DayQuil®  
Cold & Flu and Vicks® NyQuil® Cold & Flu.



## DAYTIME NIGHTTIME MULTI SYMPTOM COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59779-567
--------------	----------------	--------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-567-02	1 in 1 CARTON; Type 0: Not a Combination Product	07/27/2012	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

### Part 1 of 2

## NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

### Product Information

Item Code (Source)	NDC:59779-335
--------------------	---------------

Route of Administration	ORAL
-------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	

<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	

### Product Characteristics

<b>Color</b>	GREEN (clear, bright green)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	FRUIT (anise / cooling menthol aroma)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-335-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/16/2011	

## Part 2 of 2

### DAYTIME NON DROWSY

acetaminophen, dextromethorphan hbr, phenylephrine hcl solution

### Product Information

<b>Item Code (Source)</b>	NDC:59779-656
<b>Route of Administration</b>	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH)	DEXTROMETHORPHAN	10 mg

(DEXTROMETHORPHAN - UNII: 7355X3ROTS)	HYDROBROMIDE	in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>BUTYLATED HYDROXYANISOLE</b> (UNII: REK4960K2U)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A)	
<b>SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM</b> (UNII: 3980JIH2SW)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

## Product Characteristics

<b>Color</b>	ORANGE (clear)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MENTHOL (strong menthol, weak fruit)	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-656-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	06/09/2006	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/27/2012	

**Labeler -** CVS Pharmacy (062312574)

