

COLD AND FLU DAY- acetaminophen, dextromethorphan hbr, phenylephrine hcl solution
Walgreen Company

Walgreens 44-012

Active ingredients (in each 15 mL)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - nasal congestion
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
 - headache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user has

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- heart disease
- diabetes
- thyroid disease
- liver disease
- high blood pressure

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

Other information

- **each 15 mL contains:** sodium 13 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C yellow #6, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose

Questions or comments?

1-800-426-9391

Principal display panel

NDC 0363-9912-02

Walgreens

WALGREENS PHARMACIST RECOMMENDED†

Compare to the active ingredients
in Vicks® DayQuil® Cold & Flu††

DAY

Cold & Flu

ACETAMINOPHEN/

PAIN RELIEVER / FEVER REDUCER

DEXTROMETHORPHAN HBr /

COUGH SUPPRESSANT

PHENYLEPHRINE HCl /

NASAL DECONGESTANT

- Relieves headaches, fever, sore throat, minor aches & pains, cough

- Nasal congestion

Ages 6 Years and Over

Menthol
flavor

12 FL OZ (355 mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED
NECK WRAP IS BROKEN OR MISSING**

PARENTS:

**Learn about teen medicine abuse
www.StopMedicineAbuse.org**

**Do Not Take
Daytime and
Nighttime
Products at the
Same Time.**

†Our pharmacists recommend the Walgreens brand.

We invite you to compare to national brands.

††This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil® Cold & Flu.

DISTRIBUTED BY: **WALGREEN CO.**
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2022 Walgreen Co.

50844 REV0323A01202

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- heart disease
- diabetes
- thyroid disease
- liver disease
- high blood pressure

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

¹Our pharmacists recommend the Walgreens brand.
We invite you to compare to national brands.
^{††}This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil® Cold & Flu.
DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2022 Walgreen Co.

ITEM 852225 W00000-0000-0



3 11917 12596 1

W3ORG0922-F
50844 REV0323A01202 REV0423

**No Print / No Varnish Area
Lot # and Exp. Info**

PEEL HERE FOR MORE DRUG FACTS

NDC 0363-9912-02



Compare to the active ingredients in Vicks® DayQuil® Cold & Flu^{††}

DAY

Cold & Flu

**ACETAMINOPHEN /
PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr /
COUGH SUPPRESSANT
PHENYLEPHRINE HCl /
NASAL DECONGESTANT**

- Relieves headaches, fever, sore throat, minor aches & pains, cough
- Nasal congestion

Ages 6 Years and Over



Menthol flavor

12 FL OZ (355 mL)

Drug Facts

TAMPER EVIDENT: DO NOT USE IF PRINTED NECK WRAP IS BROKEN OR MISSING

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
- sore throat
- nasal congestion
- minor aches and pains
- cough due to minor throat and bronchial irritation
- headache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user has

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed

- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

Other information

- each 15 mL contains: sodium 13 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients anhydrous citric acid, FD&C yellow #6, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose

Questions or comments? 1-800-426-9391

Walgreens 44-012

COLD AND FLU DAY

acetaminophen, dextromethorphan hbr, phenylephrine hcl solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:0363-9912

Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg in 15 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL	
Inactive Ingredients				
Ingredient Name			Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
GLYCERIN (UNII: PDC6A3C0OX)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)				
SODIUM METABISULFITE (UNII: 4VON5FNS3C)				
SACCHARIN SODIUM (UNII: SB8ZUX40TY)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
Product Characteristics				
Color	orange	Score		
Shape		Size		
Flavor	MENTHOL	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9912-02	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/14/2022	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M012	11/14/2022	

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(0363-9912) , pack(0363-9912)

Revised: 11/2023

Walgreen Company