MECLIZINE HCL 12.5 MG- meclizine hcl 12.5 mg tablet Advance Pharmaceutical Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Meclizine HCL 12.5 caplets

Drug Facts

Active indredient (in each caplet)

Meclizine HCL 12.5 mg

Purpose

Antiemetic

Uses

- prevents and treats nausea, vomiting, or dizziness due to motion sickness
- for other uses, consult your doctor

Warnings

Ask a docotor before use if you have

- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- you may get drowsy
- avoid alcoholic drinks
- alcohol, sedatives and tranquillizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Ask a doctor before use if you have

- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

When using this prodcut

- you may get drowsy
- avoid alcoholic drinks
- alcohol, sedatives and tranqulizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults & children 12 years and over: 2-4 caplets once daily
- children under 12 years: ask a doctor

Other information

- each caplet contains: calcium 51 mg
- store at room temperature 15°-30°C (59°-86°F)

Inactive ingredients

croscarmellose sodium, dicalcium phosphate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?

call 631-981-4600, 8.30 am-4.30 pm ET, Monday - Friday


TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR

MISSING

NDC: 17714-117-01

Meclizine 12.5mg caplets





MECLIZINE HCL 12.5 MG

meclizine hcl 12.5 mg tablet

Product Type HUMAN OTC DRUG Item Code (Source) NDC:17714-117

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

MECLIZINE HYDRO CHLO RIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570) MECLIZINE HYDRO CHLO RIDE 12.5 mg

Inactive Ingredients

Ingredient Name Strength

SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)

CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)	
CALCIUM PHO SPHATE, DIBASIC, ANHYDRO US (UNII: L11K75P92J)	

Product Characteristics				
Color	white	Score	2 pieces	
Shape	OVAL	Size	13mm	
Flavor		Imprint Code	AP;117	
Contains				

ı	Packaging				
ı	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
ı	1 N	DC:17714-117-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	07/01/2013	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part336	07/01/2013		

Labeler - Advance Pharmaceutical Inc. (078301063)

Registrant - Advance Pharmaceutical Inc. (078301063)

Establishment				
Name	Address	ID/FEI	Business Operations	
Advance Pharmaceutical Inc.		078301063	manufacture(17714-117)	

Revised: 10/2017 Advance Pharmaceutical Inc.