

HUMCO MILK OF MAGNESIA- magnesium hydroxide suspension
Humco Holding Group, Inc.

Humco Milk of Magnesia USP

Drug Facts

Active ingredient

Magnesium Hydroxide 400 mg per teaspoon (5 mL)

Purpose

Antacid/Saline Laxative

Use

As an antacid: helps relieve acid ingestion, heartburn and sour stomach.

As a laxative: For the temporary relief of occasional constipation. Bowel movement is generally produced in 1.2 to 6 hours.

WARNINGS

Ask a doctor before use if you have.

Abdominal pain, nausea, vomiting, kidney disease, are on a magnesium restricted diet, a sudden change in bowel habits lasting longer than 14 days, if you are taking prescription drugs. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor.

If you have rectal bleeding or have no bowel movement after using this product. this could be signs of a serious condition, if you need to use a laxative for more than one week. Frequent or prolonged use may result in the dependence on laxatives.

When using this product.

As an antacid, do not take more than the maximum recommended daily dose in a 24 hr. period or take the maximum dose for more than two weeks, except under the advice of a doctor. May have laxative effect.

If pregnant or breastfeeding ask a health professional before use.

Keep out of reach of children.

In the case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Directions (Shake well before use)

As an antacid: Adults and children 12 yrs. of age and older: 1 to 3 teaspoonfuls with a little water, or as directed by a doctor.

As a laxative: Adults and children 12 yrs. of age and older: 1 to 4 tablespoonfuls followed by a full 8 oz. glass of water.

Children 6 thru 11 yr. of age: 1 to 2 tablespoonfuls followed by a full 8 oz. glass of water.

children 2 thru 5 yrs. of age: 1 to 3 teaspoonfuls followed by a full 8 oz. glass of water.

Children under 2 yrs. of age: consult a doctor before use.

Other information

acid neutralizing capacity: 10.0 mEq per teaspoon.

keep tightly closed and avoid freezing.

Inactive ingredient.

Purified water

Principal display panel

Drug Facts	
Active ingredient Magnesium Hydroxide 400 mg per teaspoonful (5 mL)	Purpose Antacid/Saline laxative
Indications ■ As an antacid for the relief of acid indigestion, heartburn and sour stomach. ■ As a laxative For the relief of occasional constipation. Bowel movement is generally produced in 1/2 to 6 hours.	
Warnings Ask a doctor before use if you have ■ abdominal pain ■ nausea ■ vomiting ■ kidney disease ■ are on a magnesium restricted diet ■ a sudden change in bowel habits lasting longer than 14 days. ■ if you are taking prescription drugs. Antacids may interact with certain prescription drugs. ■ Stop use and ask a doctor if you have rectal bleeding or have no bowel movement after using this product. These could be signs of a serious condition. ■ if you need to use a laxative for more than 1 week. Frequent or prolonged use may result in the dependence on laxatives.	
When using this product ■ as antacid, do not take more than the maximum recommended daily dose in a 24 hr. period or take the maximum dose for more than two weeks, except under the advice of a doctor. May have laxative effect.	
If pregnant or breast feeding ask a health professional before use. Keep out of reach of children.	

**MILK OF
MAGNESIA**
SALINE LAXATIVE • ANTACID

HUMCO
A trusted name
in home health care

* Safety Sealed
**Gastro-
intestinal**

USP

• ALL
NATURAL
STIMULANT
LAXATIVE

• ANTACID

16 FL. oz
(473 mL)

WARNING: Do not use if Tamper Evident Seal is broken or missing. This product is sealed with a breakaway cap ring.

Drug Facts (CONTINUED)	
Directions (Shake well before use) ■ As an antacid: Adults and children 12 yrs. of age and older: 15 to 45 milliliters every 4 hours with a little water, or as directed by a doctor. ■ As a laxative: Adults and children 12 yrs. of age and older: 30 to 60 milliliters followed by a full 8 oz. glass of water. ■ Children 6 thru 11 yrs. of age: 15 to 30 milliliters followed by a full 8 oz. glass of water. ■ Children 2 thru 5 yrs. of age: 5 to 15 milliliters followed by a full 8 oz. glass of water. ■ Children under 2 yrs. of age: consult a doctor before use.	
Other information ■ acid neutralizing capacity: 10.0 mEq. per teaspoon. ■ keep tightly closed and avoid freezing.	
Inactive ingredients Purified Water, Sodium Hypochlorite, Xanthan Gum.	

R100917

HUMCO, TexasKana, TX 75501
Questions or Comments? ☎1-800-662-3435
"We Help People Feel Better"
Go to: www.simplehomemedies.com to find helpful hints and learn more about Milk of Magnesia.

HUMCO MILK OF MAGNESIA

magnesium hydroxide suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0395-1672
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	80 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0395-1672-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/31/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	01/31/2019	

Labeler - Humco Holding Group, Inc. (825672884)

Registrant - Pharma Nobis, LLC (118564114)

Establishment

Name	Address	ID/FEI	Business Operations
Pharma Nobis, LLC		118564114	manufacture(0395-1672) , analysis(0395-1672) , pack(0395-1672) , label(0395-1672)

Revised: 12/2023

Humco Holding Group, Inc.